arXiv:1612.00757v1 [math.CV] 2 Dec 2016

TWISTED HODGE FILTRATION: CURVATURE OF THE DETERMINANT

PHILIPP NAUMANN

Abstract. Given a holomorphic family $f : \mathcal{X} \to S$ of compact complex manifolds and a relative ample line bundle $L \to \mathcal{X}$, the higher direct images $R^{n-p}f_*\Omega^p_{\mathcal{X}/S}(L)$ carry a natural hermitian metric. An explicit formula for the curvature tensor of these direct images is given in [Na16]. We prove that the determinant of the twisted Hodge filtration $F^p_L = \oplus_{i\geq p} R^{n-i}\Omega^i_{\mathcal{X}/S}(L)$ is (semi-) positive on the base S if L itself is (semi-) positive on $\mathcal{X}$.

1. introduction

For an ample line bundle L on a compact complex manifolds X of dimension n, the cohomology groups $H^{n-p}(X, \Omega^p_X(L))$ are critical with respect to Kodaira-Nakano vanishing. More generally, we consider the higher direct image sheaves $R^{n-p}f_*\Omega^p_{\mathcal{X}/S}(L)$ for a proper holomorphic submersion $f : \mathcal{X} \to S$ of complex manifolds and a line bundle $L \to \mathcal{X}$, which is positive along the fibers $X_s = f^{-1}(s)$. The understanding of this situation has applications to moduli problems. An explicit curvature formula for these higher direct images is given in [Na16]. In general, the curvature of the intermediate higher direct images contains positive and negative contributions. Therefore, we consider the twisted Hodge filtration

$$F^p_L := \oplus_{i\geq p} R^{n-i}\Omega^i_{\mathcal{X}/S}(L)$$

This allows us to offset the negative terms by the positive ones. The result is that $\det(F^p_L)$ is (semi-) positive on the base S if L is (semi-) positive and relatively ample on $\mathcal{X}$. This idea comes from [Sch16].

2. Differential geometric setup and proof of the result

First we recall the setting from [Na16]. Let $f : \mathcal{X} \to S$ be a proper holomorphic submersion and (L, h) a line bundle on $\mathcal{X}$. The curvature form of the hermitian line bundle is given by

$$\omega_{\mathcal{X}} := 2\pi \cdot c_1(L, h) = -\sqrt{-1}\partial\bar{\partial}\log h.$$

We consider the case where the hermitian bundle (L, h) is relative ample, which means that

$$\omega_{X_s} := \omega|_{X_s}$$

are Kähler forms on the n-dimensional fibers X_s. Then one has the notion of a horizontal lift v_s of a tangent vector ∂_s on the base S and we get a representative of the Kodaira-Spencer class

$$A_s := \bar{\partial}(v_s)|_{X_s}.$$

Note that we have $L_{v_s}(\omega_{X_s})^n = 0$ (see [Be11]). Furthermore, one sets

$$\varphi := \langle v_s, v_s\rangle_{\omega_{\mathcal{X}}},$$

which is called the geodesic curvature. The coherent sheaf $R^{n-p}f_*\Omega^p_{\mathcal{X}/S}(L)$ is locally free on S outside a proper subvariety. In the case $n = p$ and L ample, the sheaf $f_*(K_{\mathcal{X}/S} \otimes L)$ is locally free by the Ohsawa-Takegoshi extension theorem (see [Be09]). We assume the local freeness of

$$R^{n-p}f_*\Omega^p_{\mathcal{X}/S}(L)$$

2000 *Mathematics Subject Classification.* 32L10, 32G05, 14Dxx.

Key words and phrases. Curvature of higher direct image sheaves, Deformations of complex structures, Families, Fibrations, Twisted Hodge filtration.

in the general case, hence the statement of the Grothendieck-Grauert comparison theorem holds. Now Lemma 2 of [Sch12] applies, which says that we can represent local sections of $R^{n-p}f_*\Omega^p_{\mathcal{X}/S}(L)$ by $\bar\partial$-closed $(p,n-p)$-forms on the total space, whose restrictions to the fibers are harmonic. Let $\{\psi^1,\dots,\psi^r\}$ be a local frame of the direct image consisting of such sections around a fixed point $s\in S$. We denote by $\{(\partial/\partial s_i)|i=1,\dots,\dim S\}$ a basis of the complex tangent space T_sS of S over $\mathbb{C}$, where s_i are local holomorphic coordinates on S. The natural inner product is given by

$$(\psi^k,\psi^l)=\int_X \psi^k_{A_p\overline{B}_q}\psi^{\overline{l}}_{C_q\overline{D}_p}g^{\overline{D}_pA_p}g^{\overline{B}_qC_q}h\,g\,dV,$$

Let $A^\alpha_{i\overline{\beta}}(z,s)\partial_\alpha dz^{\overline{\beta}}=\bar\partial(v_i)_{X_s}$ be the $\bar\partial$-closed representative of the Kodaira-Spencer class of ∂_i described above. Then the cup product together with contraction defines

$$A^\alpha_{i\overline{\beta}}\partial_\alpha dz^{\overline{\beta}}\cup\ :\mathcal{A}^{0,n-p}(X_s,\Omega^p_{X_s}(L|_{X_s}))\to\mathcal{A}^{0,n-p+1}(X_s,\Omega^{p-1}_{X_s}(L|_{X_s})) \tag{1}$$

$$A^{\overline{\beta}}_{\overline{j}\alpha}\partial_{\overline{\beta}}dz^\alpha\cup\ :\mathcal{A}^{0,n-p}(X_s,\Omega^p_{X_s}(L|_{X_s}))\to\mathcal{A}^{0,n-p-1}(X_s,\Omega^{p+1}_{X_s}(L_{X_s})) \tag{2}$$

where $p>0$ in (1) and $p<n$ in (2). Note that this is a formal analogy to the derivative of the period map in the classical case (see [Gr70]). We will apply the above cup products to harmonic $(p,n-p)$-forms. In general, the results are not harmonic.

When applying the Laplace operator to (p,q)-forms with values in L on the fibers X_s, we have

$$\Box_\partial-\Box_{\bar\partial}=(n-p-q)\cdot\mathrm{id} \tag{3}$$

due to the definition $\omega_{X_s}=\omega_{\mathcal{X}}|_{X_s}$ and the Bochner-Kodaira-Nakano identity. Thus, we write $\Box=\Box_\partial=\Box_{\bar\partial}$ in the case $q=n-p$. By considering an eigenfunction decomposition and using the identity (3), we obtain that all eigenvalues of $\Box$ are 0 or greater than 1, hence the operator $(\Box-1)^{-1}$ exists. We use the notation $\psi^{\overline{l}}:=\overline{\psi^l}$ for sections ψ^l and write $g\,dV=\omega_{X_s}/n!$. The result from [Na16] is

Theorem 1. *Let $f:\mathcal{X}\to S$ be a proper holomorphic submersion and $(L,h)\to\mathcal{X}$ a relative ample line bundle. With the objects described above, the curvature of $R^{n-p}f_*\Omega^p_{\mathcal{X}/S}(L)$ is given by*

$$\begin{aligned}R^{\overline{l}k}_{i\overline{j}}(s)=&\int_{X_s}\varphi_{i\overline{j}}\cdot(\psi^k\cdot\psi^{\overline{l}})\,g\,dV\\ &+\int_{X_s}(\Box+1)^{-1}(A_i\cup\psi^k)\cdot(A_{\overline{j}}\cup\psi^{\overline{l}})\,g\,dV\\ &+\int_{X_s}(\Box-1)^{-1}(A_i\cup\psi^{\overline{l}})\cdot(A_{\overline{j}}\cup\psi^k)\,g\,dV\end{aligned}$$

If $L\to\mathcal{X}$ is ample, the only contribution, which may be negative, originates from the harmonic parts in the third term

$$-\int_{X_s}H(A_I\cup\psi^{\overline{l}})\overline{H(A_j\cup\psi^{\overline{k}})}\,g\,dV.$$

From now on, we assume that the base space S is one dimensional. Let ∂_s be a tangent vector and $A=A_s$ the corresponding $\bar\partial$-closed Dolbeault representative of the Kodaira-Spencer class. By considering an eigenvector decomposition, we have the following estimate for the curvature of $R^{n-p}\Omega^p_{\mathcal{X}/S}(L)$ for a semi-positive, relatively ample line bundle L (see also [Sch16, Corollary 5]):

Corollary 1.

$$R(A,\overline{A},\psi,\overline{\psi})\geq||H(A\cup\psi)||^2-||H(\overline{A}\cup\psi)||^2 \tag{4}$$

Now we formalize the situation: The cup products (1) and (2) give rise to cup products for cohomology classes (ψ is assumed to be harmonic).

$$A_p\ :H^{n-p}(X_s,\Omega^p_{X_s}(L|_{X_s}))\to H^{n-p+1}(X_s,\Omega^{p-1}_{X_s}(L|_{X_s}))\quad;\quad[\psi]\mapsto[H(A\cup\psi)] \tag{5}$$

$$\overline{A}_p\ :H^{n-p}(X_s,\Omega^p_{X_s}(L|_{X_s}))\to H^{n-p-1}(X_s,\Omega^{p+1}_{X_s}(L_{X_s}))\quad;\quad[\psi]\mapsto[H(\overline{A}\cup\psi)] \tag{6}$$

We have ([Sch16, Lemma 5])

Lemma 1. *$\overline{A}_p = (A_{p+1})^*$, i.e. the map $\overline{A}_p$ is the adjoint of the map A_{p+1}.*

Proof. For any harmonic $\chi \in \mathcal{A}^{(p,n-p)}(X_s, L|_{X_s})$ we have $(H(\overline{A} \cup \psi), \chi) = (\overline{A} \cup \psi, \chi) = (\psi, A \cup \chi) = (\psi, H(A \cup \chi))$. □

In the spirit of the Hodge filtration we define for $0 \leq p \leq n$

Definition 1.

$$F_L^p := \bigoplus_{i \geq p} R^{n-i}\Omega^i_{\mathcal{X}/S}(L)$$

Then we have

Theorem 2. *The line bundle $\det(F_L^p)$ is (semi-) positive on S, if L is (semi-) positive and relatively ample on $\mathcal{X}$.*

Proof. We choose a local holomorphic frame $\{\psi^\alpha\} \subset \mathcal{A}^{(p,n-p)}(X_s, L|_{X_s})$ of harmonic forms and compute

$$\begin{aligned}
\sum_\alpha ||H(A \cup \psi^\alpha)||^2 &= \sum (H(A \cup \psi^\alpha), H(A \cup \psi^\alpha)) \\
&= \sum (H(\overline{A} \cup A \cup \psi^\alpha), \psi^\alpha) \\
&= \sum ((A_p^* \circ A_p)\psi^\alpha, \psi^\alpha) \\
&= \operatorname{tr}(A_p^* A_p)
\end{aligned}$$

Analogously

$$\begin{aligned}
\sum_\alpha ||H(\overline{A} \cup \psi^\alpha)||^2 &= \sum (H(\overline{A} \cup \psi^\alpha), H(\overline{A} \cup \psi^\alpha)) \\
&= \sum (H(A \cup \overline{A} \cup \psi^\alpha), \psi^\alpha) \\
&= \sum ((\overline{A}_p^* \circ \overline{A}_p)\psi^\alpha, \psi^\alpha) \\
&= \operatorname{tr}(\overline{A}_p^* \overline{A}_p) \\
&= \operatorname{tr}(A_{p+1} A_{p+1}^*),
\end{aligned}$$

because of Lemma 1. Note that we have $\operatorname{tr}(A_p^* A_p) = \operatorname{tr}(A_p A_p^*)$. Now the result follows from Corollary 1, since the curvature of the determinant of a bundle is the trace of the curvature form of the bundle. □

References

[Be09] B. Berndtsson: *Curvature of vector bundles associated to holomorphic fibrations*, Ann. Math. **169**, 531-560 (2009).

[Be11] B. Berndtsson: *Strict and nonstrict positivity of direct image bundles*, Math. Zeitschrift, Vol. **269**, 1201-1218 (2011).

[De12] J.-P. Demailly: *Complex Analytic and Differential Geometry*, https://www-fourier.ujf-grenoble.fr/ demailly/manuscripts/agbook.pdf, Grenoble (2012).

[Gr70] P.A. Griffiths: *Periods of integrals on algebraic manifolds III: Some differential-geometric properties of the period mapping*, Publ. Math. IHES **38**, 125-180 (1970).

[MT08] C. Mourougane, S. Takayama: *Hodge metrics and the curvature of higher direct images*, Ann. Sci. Ec. Norm. Supér. (4) **41**, 905-924 (2008).

[Na16] P. Naumann: *Curvature of higher direct images*, arXiv:1611.09117 (2016).

[Sch93] G. Schumacher: *The curvature of the Petersson-Weil metric on the moduli space of Kähler-Einstein manifolds*, Ancona, V. (ed.) et al., Complex analysis and geometry. New York: Plenum Press. The University Series in Math., 339-354 (1993).

[Sch12] G. Schumacher: *Positivity of relative canonical bundles and applications*, Invent. Math. **190**, 1-56 (2012).

[Sch13] G. Schumacher: *Erratum to: Positivity of relative canonical bundles and applications*, Invent. Math. **192**, 253-255 (2013).

[Sch16] G. Schumacher: *Moduli of canonically polarized manifolds, higher order Kodaira-Spencer maps, and an analogy to Calabi-Yau manifolds*, to appear in (2016).

Fachbereich Mathematik und Informatik, Philipps-Universität Marburg, Lahnberge, Hans-Meerwein-Strasse, D-35032 Marburg,Germany

E-mail address: naumann@mathematik.uni-marburg.de